

Film Canister Rocket Instructions



1. Put on safety goggles or some kind of eye protection and head outside – no really, when this works, that film canister really flies! If you want to try the indoor version, do not turn the canister upside down in step 5.
2. Break the antacid tablet in half.
3. Remove the lid from the film canister and put a teaspoon (5 ml) of water into the canister. Do the next 2 steps quickly
4. Drop the tablet half into the canister and snap the cap onto the canister (make sure that it snaps on tightly.)
5. Quickly put the canister on the ground CAP SIDE DOWN and STEP BACK at least 2 meters.
6. About 10 seconds later, you will hear a POP! and the film canister will launch into the air!

Caution: If it does not launch, wait at least 30 second before examining the canister. Usually the cap is not on tight enough and the build up of gas leaked out.

Film Canister Rocket Instructions



1. Put on safety goggles or some kind of eye protection and head outside – no really, when this works, that film canister really flies! If you want to try the indoor version, do not turn the canister upside down in step 5.
2. Break the antacid tablet in half.
3. Remove the lid from the film canister and put a teaspoon (5 ml) of water into the canister. Do the next 2 steps quickly
4. Drop the tablet half into the canister and snap the cap onto the canister (make sure that it snaps on tightly.)
5. Quickly put the canister on the ground CAP SIDE DOWN and STEP BACK at least 2 meters.
6. About 10 seconds later, you will hear a POP! and the film canister will launch into the air!

Caution: If it does not launch, wait at least 30 second before examining the canister. Usually the cap is not on tight enough and the build up of gas leaked out.